

Non-linear charge reduction effect in strongly-coupled plasmas

D. Sarmah^{1,2,3}, M. Tessarotto^{1,3} and M. Salimullah⁴

¹*Department of Mathematics and Informatics,*

University of Trieste, Trieste, Italy

² *International Center for Theoretical Physics, ICTP/TRIL*

Program, Trieste, Italy

³ *Consortium for Magnetofluid Dynamics, Trieste, Italy*

⁴ *Department of Physics,*

Jahangirnagar University, Savar, Dhaka,

Bangladesh

(Dated: February 2, 2008)

Abstract

The charge reduction effect, produced by the nonlinear Debye screening of high-Z charges occurring in strongly-coupled plasmas, is investigated. An analytic asymptotic expression is obtained for the charge reduction factor (f_c) which determines the Debye-Hückel potential generated by a charged test particle. Its relevant parametric dependencies are analyzed and shown to predict a strong charge reduction effect in strongly-coupled plasmas.

PACS numbers: 51.50+v, 52.20-j, 52.27.Gr

A basic aspect of plasma physics is the so-called Debye shielding of the electrostatic potential. This consists in the property of plasmas (or electrolytes [1]), either quasi-neutral or non-neutral, to shield the electrostatic field produced by charged particles (to be also denoted as test particles) immersed in the same system. This result has fundamental consequences on plasma phenomenology, since it actually limits the range of static Coulomb interactions inside the Debye sphere, i.e., at a distance $\rho \leq \lambda_D$ from the test particle, λ_D being the Debye length. As usual here $\lambda_D \equiv \left(\sum_s \lambda_{Ds}^{-1} \right)^{-1}$, where the sum is carried out on all plasma species and $\lambda_{Ds} = \sqrt{\frac{T_s}{4\pi Z_s^2 e^2 N_{os}}}$, T_s and N_{os} being respectively the s -species temperature and number density (the latter defined in the absence of test particles). In fact, when both the particles and the plasma are assumed non-relativistic, small-amplitude, stationary (or slowly time- and space-varying), electrostatic perturbations generated by isolated test particles, result effectively shielded in the external domain, i.e., at distances larger than the Debye length λ_D . The renewed interest in this problem is particularly related to dusty plasmas or colloidal suspensions [2, 3] which are characterized by the presence of a large fraction of highly charged particles (grains), i.e., having an electric charge $Z_d e$ with $|Z_d| \gg 1$. It is well established that the phenomenon of Debye shielding of the electrostatic potential generated by a slowly moving or stationary charged test particle, when occurring in strongly coupled ionized gases, manifests peculiar properties. These are produced, as a consequence of the nonlinear plasma response occurring at distances smaller than the Debye length λ_D , when the electrostatic potential (Φ) results locally such that $\hat{\Phi} \equiv |e\Phi|/T_o \sim 1$ (or even $\gg 1$), $\hat{\Phi}$ being the normalized electrostatic potential evaluated at a suitable characteristic distance ρ_o from the position of the test particle $\mathbf{r}(t)$ and $T_o \equiv T_i$. The distance ρ_o can be identified with the radius of the local plasma sheath around a test particle. For spherically-symmetric test particles such a domain can be identified with the spherical subset of the configuration space, of radius $\rho_o \equiv \left(\frac{3}{4\pi N_o} \right)^{1/3}$ (average interparticle distance), $N_o \equiv N_{oi}$ being the density of the ion species, in which each test particle can be considered as isolated. Ionized gases (or electrolytes) can be classified according to the characteristic dimensionless parameters $x_o = \rho_o/\lambda_D \equiv g^{1/3}$ and $\Gamma \equiv \beta/x_o$ (where $\beta = \frac{Z_d e^2}{4\pi T_o \lambda_D}$ is denoted as the dimensionless electric charge of the test particle), g and Γ being respectively so-called *plasma* and *Coulomb coupling parameters*. In particular, for plasmas the high-density requirement $x_o^3 \equiv g \ll 1$ is satisfied by assumption and can be interpreted as the condition that the total number of

mutually interacting particles in a Debye sphere (expressed by $1/g$) results $\gg 1$. Instead, the ordering of parameter Γ remains in principle arbitrary. Thus, the orderings $\Gamma \ll 1$ and $\Gamma \sim 1$ (or $\Gamma \gg 1$) correspond respectively to so-called weakly and strongly-coupled plasmas. Dusty and colloidal plasmas may be characterized by Z_d up to $10^4 - 10^6$ with typical grain size smaller than 10^{-4} cm and with plasma temperature and the ion density $T_o \sim 1$ eV, $N_o \sim 10^8 - 10^{10}$ cm $^{-3}$. In this case the Coulomb parameter for a negatively-charged grain in the presence of the plasma sheath produced by hydrogen ions can result as large as $\Gamma \cong (10^{-2} - 10^{-3})Z_d$, while the dimensionless radius of the ion plasma sheath can be estimated $x_o \cong 0.01 - 0.05$. Therefore, dusty plasmas typically result strongly coupled if $Z_d \gtrsim 10^3$. For these plasmas it is important to be able to evaluate the electrostatic potential generated by dust grains which is expected to be strongly influenced by Debye shielding. It is well-known that, in general, this phenomenon occurs provided suitable physical assumptions are introduced. In particular, the plasma must be assumed appropriately close to kinetic Maxwellian equilibrium, in which each particle species is described by a Maxwellian kinetic distribution function carrying finite fluid fields [defined respectively by the number density, temperature and flow velocity $(N, T, \mathbf{V})$]. In the absence of test particles these fluid fields must be assumed slowly varying in a suitable sense, or constant, both with respect to position ($\mathbf{r}$) and time (t). In this regard it is important to remark that the appropriate treatment of the plasma sheath surrounding each test particle is essential also for the validity of the mathematical model for the Debye screening problem, i.e., for the existence of classical solutions of the Debye screening problem, which do not exist when letting $x_o = 0$ [4]. As a consequence the electrostatic potential depends necessarily on the dimensionless parameter x_o . In particular, in the domain $x \geq x_o$ the dimensionless potential $\widehat{\Phi}_{x_o}(x)$ generated by a dust grain of charge $Z_de < 0$, fulfilling the boundary conditions $\left. \frac{d\widehat{\Phi}_{x_o}(x)}{dx} \right|_{x=x_o} = -\frac{\Gamma}{x_o}$ and $\lim_{x \rightarrow \infty} \widehat{\Phi}_{x_o}(x) = 0$, can be proven to satisfy the *integral Debye-Poisson equation*

$$\widehat{\Phi}_{x_o}(x) = \frac{\beta}{x} - \left[\frac{1}{x} \int_{x_o}^x dx' x'^2 + \int_x^\infty dx' x' \right] S(x', x_o), \quad (1)$$

yielding for $x = x_o$ the constraint equation

$$\widehat{\Phi}_o = \Gamma - \int_{x_o}^\infty dx' x' S(x', x_o). \quad (2)$$

Here $\widehat{\Phi}_o \equiv \widehat{\Phi}_{x_o}(x_o)$, while in the case of a negatively-charged test particles in a quasi-neutral three species (electron, ion and dust) plasma the source term $S(x, x_o)$ reads $S(x, x_o) \equiv \Theta(x -$

x_o) $\left[\xi_i \exp \left\{ |Z_i| \widehat{\Phi}_{x_o}(x) \right\} - \xi_e \exp \left\{ -\widehat{\Phi}_{x_o}(x) \right\} - \xi_d \exp \left\{ -|Z_d| \widehat{\Phi}_{x_o}(x) \right\} \right]$, $\Theta(x-x_o)$ being the Heaviside function. Moreover, ξ_s is defined by the ratio $\xi_s = \frac{1}{|Z_s|} \frac{\lambda_D^2}{\lambda_{Ds}^2}$. In the case in which the contribution due to the dusty species is negligible, and moreover $|Z_i| = 1$, $T_i = T_e$, the previous expression reduces to the customary value $S(x, x_o) \equiv \Theta(x - x_o) \sinh \widehat{\Phi}_{x_o}(x)$. Another useful representation of Eq.(1) can also be obtained by means of the transformation

$$\widehat{\Phi}_{x_o}(x) = \widehat{\Phi}_o \exp \{y(x, x_o)\}, \quad (3)$$

and there results, by consistency with the integral DP equation (1), $y(x_o, x_o) = 0$ and $y'(x, x_o)|_{x=x_o} = -\alpha$, with $\alpha = \Gamma/x_o \widehat{\Phi}_o$. This delivers for $y(x, x_o)$ the integral equation

$$y(x, x_o) = y_o(x, x_o) + F(x, x_o), \quad (4)$$

$$y_o(x, x_o) = -\alpha(x - x_o), \quad (5)$$

where $F(x, x_o)$ is the solution of the integral equation

$$F(x, x_o) = \int_{x_o}^x dx' (x - x') \left\{ S(x', x_o) - y(x', x_o)^2 - \frac{2}{x'} y'(x', x_o) \right\}, \quad (6)$$

with $S(x', x_o)$ to be expressed in terms of $y(x', x_o)$ by means of Eq.(3). On the other hand, for finite x_o , the Debye-Hückel asymptotic approximation for the electrostatic potential can be proven to hold when measuring the potential at a position $\mathbf{r}$ sufficiently far from the position of the test particle $\mathbf{r}(t)$. In fact this involves imposing that the so-called *weak-field condition* $\widehat{\Phi}_{x_o}(x) \ll 1$ be locally satisfied. As a consequence, the normalized electrostatic potential generated by a spherically-symmetric point particle of charge Ze can be approximated by

$$\widehat{\Phi}_{x_o}(x) \cong \widehat{\Phi}_{x_o}^{(ext)}(x) \equiv \frac{q}{4\pi x} e^{-\Delta x}, \quad (7)$$

where $\Delta x \equiv x - x_o$, $q = q(x_o, \beta)$ is a suitable dimensionless effective electric charge and $\widehat{\Phi}_{x_o}^{(ext)}(x)$ is usually known as DH potential. The latter identifies the *external asymptotic solution* which holds in the subset of the external domain, outside the Debye sphere, in which the weak-field condition is satisfied. The remaining notation is standard, thus $\rho = |\mathbf{r} - \mathbf{r}(t)|$ is the distance from the point charge and $x \equiv \rho/\lambda_D \in [x_o, \infty]$ is the corresponding normalized distance. The weak-field condition is manifestly locally satisfied both for strongly and weakly coupled plasmas, provided there results $x > 1$, with x suitably large (i.e., $x \gg 1$). However, in the case in which there results either $x_o \ll 1$ and/or $\Gamma \gg 1$, i.e., for strongly-coupled

plasmas, the electrostatic potential may actually decay on a scale much shorter than the Debye length and precisely on a scale suitably close to the boundary of the plasma sheath, i.e., for $x_o \lesssim x \ll 1$. This conclusion is consistent both with previous analytic estimates based on the exact solution of the 1D Debye screening problem [5] and direct numerical simulations of the 3D Debye screening problem [6, 7, 8, 9]. It implies, however, that the DH approximation typically holds also at distances suitably larger than the boundary of the plasma sheath x_o and which extend also to values comparable to the Debye length, i.e., for $x_o \ll x \lesssim 1$ [10]. Finally, as a further consequence, the effective electric charge carried by the DH potential q may appear significantly reduced with respect to the case of weakly coupled plasmas [6, 8, 9, 11, 12]. To display this effect, it is convenient to represent the normalized effective charge of the test particle in the form $q \equiv \beta f_c$, f_c being the *charge reduction factor*. By dimensional analysis, it follows that f_c must be of the form $f_c = f_c(x_o, \beta, \xi_s)$, i.e., f_c is function, to be determined, of the only dimensionless parameters of the problem, namely x_o and Γ (or β) and moreover the ratios ξ_s (with $s = e, i, d$). It is well-known that for weakly-coupled plasmas q can be approximated by the normalized electric charge of the isolated test particle, i.e., $q \cong \beta$, which implies $f_c \cong 1$. Instead, for strongly-coupled plasmas, the value of charge reduction factor, determined either via one-dimensional analytic estimates [5] or by means of a variety of numerical simulation methods (see, for example, [6, 7, 8, 9]), has been found to be notably smaller than unity, thus suggesting the existence of a possible significant *charge reduction effect*. Indeed the investigation of the effective interactions characterizing high-Z grains in plasmas has attracted interest in recent years especially for their role in dusty plasmas [11, 12]. However, for strongly-coupled plasmas the precise form of the function f_c is still unknown and, in particular, an analytic estimate of the effective charge characterizing the DH potential in strongly-coupled plasmas is not yet available.

Besides being of obvious interest as a still unsolved mathematical problem, it is especially important from the physical standpoint to estimate f_c as a function of the parameters x_o and Γ for strongly coupled plasmas characterized either by a high density and/or by the presence of a large fraction of highly charged particles (grains), such as dusty plasmas and colloidal suspensions [2, 3].

In a previous work [4], the Debye screening problem (DSP) has been formulated for an electron-hydrogen plasma in order to analyze the asymptotic properties of its solutions near the plasma sheath (i.e., for x suitably close to x_o). For this purpose the case of a strongly-

coupled plasma was investigated in which the parameters x_o and Γ satisfy the asymptotic conditions, to be denoted as *strong-coupling ordering*, $\Gamma \sim \frac{1}{O(\delta)} \gg 1$ with $x_o \sim O(\delta^k)$, $k = 0, 1$ and δ an infinitesimal. The similar case of three-species plasma can be obtained directly. As a result, invoking for $\widehat{\Phi}_{x_o}(x)$ the representation (3), together with Eqs.(4),(5) and imposing the appropriate boundary conditions for $\widehat{\Phi}_{x_o}(x)$, it follows that the integral on the r.h.s. of Eq.(2) can be estimated asymptotically to yield an asymptotic equation for the initial condition $\widehat{\Phi}_o$, i.e.,

$$\widehat{\Phi}_o(x_o, \beta) \cong \frac{1}{|Z_i|} \ln \left\{ \frac{|Z_i|}{\xi_i} \frac{\Gamma}{x_o^2} [\Gamma - \widehat{\Phi}_o] \right\}. \quad (8)$$

In particular, for an electron-hydrogen plasma in which the contribution to λ_D due to the dusty species is negligible, there results $|Z_i| = 1$ and $\xi_i = 1/2$ and this reduces (8) to the expression given in Ref. [4]. In addition, neglecting higher-order infinitesimals of order $O(\delta^r)$ ($r > 0$ being a suitable real number), the following inequality can immediately be proven

$$f_c \leq c^{(a)}(x_o, \beta) [1 + O(\delta^r)], \quad (9)$$

$$c^{(a)}(x_o, \beta) \equiv x_o \widehat{\Phi}_o(x_o, \beta), \quad (10)$$

$c^{(a)}(x_o, \beta)$ being an *asymptotic upper bound* for the charge reduction factor. This result, as indicated below, is relevant to investigate the charge reduction effect in strongly-coupled plasmas.

In this paper we propose a more accurate asymptotic approximation for f_c , potentially useful to analyze the charge reduction effect of high-Z test particles taking place in strongly-coupled plasmas. In particular, we intend to prove that, in validity of the previous strong-coupling ordering, the charge reduction factor can be approximated in the form:

$$f_c(x_o, \beta) \cong c^{(b)}(x_o, \beta) \equiv x_c e^{\Delta x_c} \widehat{\Phi}_{x_o}^{(int)}(x_c) / \beta, \quad (11)$$

where, x_c is defined so that $x_c > x_o$ and is provided by the root of the equation

$$y'(x_c, x_o) = -1 - \frac{1}{x_c}. \quad (12)$$

Here $\widehat{\Phi}_{x_o}^{(int)}(x)$ denotes the leading-order contribution to the internal asymptotic solution of the Debye-Poisson equation (1), which can be obtained from the representation (3),(4),(5)

and the integral equation (6) and holds in an appropriate neighborhood of the plasma sheath (x_o) [4]. There results

$$\widehat{\Phi}_{x_o}^{(int)}(x) \cong \exp \{y_o(x, x_o)\}, \quad (13)$$

while $y'(x, x_o)$ reads

$$\begin{aligned} y'(x, x_o) \cong & -\alpha - \alpha^2 \Delta x + 2\alpha \log \frac{x}{x_o} + \\ & + \frac{\xi_i}{\widehat{\Phi}_o} \int_{x_o}^x dx' \exp \left\{ |Z_i| \widehat{\Phi}_o \exp \{-\alpha \Delta x'\} + \alpha \Delta x' + \frac{1}{2} \alpha^2 \Delta x'^2 - 2\alpha \left[x' \log \frac{x'}{x_o} - \Delta x' \right] \right\}, \end{aligned} \quad (14)$$

where $\Delta x' = x' - x_o$. Eqs.(11) and (12) uniquely follow by imposing suitable matching conditions between the internal and external asymptotic solutions, $\widehat{\Phi}_{x_o}^{(ext)}(x)$ and $\widehat{\Phi}_{x_o}^{(int)}(x)$, precisely

$$\widehat{\Phi}_{x_o}^{(int)}(x_c) = \widehat{\Phi}_{x_o}^{(ext)}(x_c), \quad (15)$$

$$\left. \frac{\partial}{\partial x} \widehat{\Phi}_{x_o}^{(int)}(x) \right|_{x=x_c} = \left. \frac{\partial}{\partial x} \widehat{\Phi}_{x_o}^{(ext)}(x) \right|_{x=x_c}. \quad (16)$$

It is immediate to prove that Eq.(16) is equivalent to the condition of stationarity for the function $f_c^{(a)}(x, x_o, \beta)$

$$\left. \frac{\partial}{\partial x} f_c^{(a)}(x, x_o, \beta) \right|_{x=x_c} = 0. \quad (17)$$

Therefore Eq.(17) also furnishes the same asymptotic approximation for the charge reduction factor given by Eq.(11). The accuracy of the asymptotic approximation $c^{(b)}(x_o, \beta)$ and of the upper bound $c^{(a)}(x_o, \beta)$ can be tested by comparing them with the estimate of the charge reduction factor obtained from direct numerical simulations of the 3D Debye screening problem. For this purpose the function

$$g(x, x_o, \beta) = x e^{x-x_o} \widehat{\Phi}_{x_o}(x) / \beta \quad (18)$$

has to be evaluated numerically by solving the Debye-Poisson equation. The charge reduction factor is defined as the limit

$$f_c(x_o, \beta) = \lim_{x \rightarrow +\infty} x e^{x-x_o} \widehat{\Phi}_{x_o}(x) / \beta. \quad (19)$$

In practice, for numerical estimates it can be approximated by $f_c(x_o, \beta) \cong g(x_f, x_o, \beta)$ with x_f to be suitably defined. It is important to stress that, in the limit $\delta \rightarrow 0$, one expects

$x_c \rightarrow 0$, $x_f \rightarrow 0$ and moreover

$$\lim_{\delta \rightarrow 0} c^{(b)}(x_o, \beta) = \lim_{\delta \rightarrow 0} f_c(x_o, \beta). \quad (20)$$

In addition, it must result also

$$\lim_{\delta \rightarrow 0} c^{(b)}(x_o, \beta) = \lim_{\delta \rightarrow 0} c^{(a)}(x_o, \beta). \quad (21)$$

Therefore, for δ small enough (i.e., for $\Gamma \gg 1$ and/or $x_o \ll 1$), it is actually possible to approximate $f_c(x_o, \beta)$ both in terms of $c^{(b)}(x_o, \beta)$ and $c^{(a)}(x_o, \beta)$. Numerical simulations indicate, however, that the asymptotic estimate $c^{(b)}(x_o, \beta)$ actually converges rapidly to $f_c(x_o, \beta)$ and furnishes therefore a more accurate approximation than the upper bound $c^{(a)}(x_o, \beta)$. Nevertheless, particularly thanks to Eq.(21), $c^{(a)}(x_o, \beta)$ can still be used for order-of-magnitude estimates.

In figures 1 the result of two numerical simulations are reported, for the case of an electron-hydrogen plasma and for $\xi_i = 1/2$, which correspond respectively to $(x_o, \beta) = (0.08, 2.5)$ and $(x_o, \beta) = (0.15, 10)$. In both cases the matching point x_c is found to be close to the boundary of the plasma sheath, which implies that the DH potential actually approximately applies for $x \geq x_c$. The upper bound estimate $c^{(a)}(x_o, \beta)$ can be used for order-of-magnitude estimates of the charge reduction effect. Finally, the parametric dependencies of $c^{(a)}(x_o, \beta)$ with respect to β and x_o are displayed for selected values of the parameters. Figures 2 demonstrate that for typical values of the Coulomb parameter Γ and the radius of the plasma sheath x_o the effective electric charge of the DH potentially appear dramatically reduced.

The present results are relevant for the investigation of strongly-coupled plasmas, such as dusty plasmas, in which a significant fraction of high- Z charged particles is present. In fact, the charge reduction effect has fundamental consequences on the Debye screening phenomenon, since it actually limits the magnitude of the Coulomb interactions produced by these particles. The estimate of the charge reduction factor f_c , here obtained, is a necessary prerequisite for the description of a variety of kinetic processes (for example, condensation [13], collisional processes [14, 15], etc.), as well as electrostatic or electromagnetic instabilities in dusty plasmas (see for example [16, 17]).

Acknowledgments

Work supported by PRIN Research Program “*Programma Cofin 2004: Modelli della teoria cinetica matematica nello studio dei sistemi complessi nelle scienze applicate*” (MIUR Italian Ministry), the ICTP/TRIL Program (ICTP, Trieste, Italy), and the Consortium for Magnetofluid Dynamics, University of Trieste, Italy.

- [1] P. Debye and E. Hückel, Phys. Zeit. **24**(9), 186 (1923); see also p.217 in *The collected papers of Peter J.W. Debye*, Ox Bow Press, Conn., USA, (1988).
- [2] P. Pieransky, Contemp.Phys **24**, 25 (1983).
- [3] H. Thomas et al., Phys. Rev.Lett. **73**, 652 (1994).
- [4] D. Sarmah, M. Tassarotto and M. Salimullah, *Plasma-sheath effects on the Debye screening problem*, <http://arxiv.org/abs/physics/0512213> (2005).
- [5] R. Clemente and P. Martin, J.Phys. Soc.J. **61**(6), 1969 (1992).
- [6] M.O. Robbins, K .Kremer, G.S. Grest, J.Chem. Phys. **88**, 3286 (1988).
- [7] G. Dupont, Mol. Phys.**79**, 453 (1996).
- [8] E. Allayrov, H. Löwen, S. Trigger, Phys. Rev.E **57**, 5518 (1998).
- [9] O. Bysternko and A. Zagorodny, Phys.Lett. **A 255**, 325 (1999).
- [10] A. Gregoratto, C. Lo Surdo, M. Tassarotto and R. Zorat, Proc. 2nd Symposium on Plasma Dynamics: Theory and Applications, Consortium for Magnetofluid Dynamics and C.D.C., Udine, Italy, p.39 (1992).
- [11] V.N. Tsytovich, Comm.Plasma Phys. **15** (6), 349 (1974).
- [12] O. Bysternko, T. Bysternko and A. Zagorodny, Cond.Matter Physics **6** (3), 425 (2003).
- [13] P.M. Bellan, Phys. Plasmas **11**, 3368 (2004).
- [14] V.N.Tsytoich and U. de Angelis, Phys. Plasma **6**, 1093 (1999).
- [15] P. Ricci, G. Lapenta, U. de Angelis and V.N. Tsytovich, Phys. Plasma **8**, 769 (2001).
- [16] M.Salimullah, M.Rosenberg, Phys. Lett. A **254**, 347 (1999).
- [17] P.K. Shukla, M. Salimullah and G. Sorasio, Phys.Plasmas **9**, 5121 (2002).




FIG. 1: Comparison between the asymptotic approximation of the charge reduction factor $c^{(a)}(x_o, \beta)$, the numerical estimate $f_c(x_o, \beta)$ and the upper bound estimate $c^{(b)}(x_o, \beta)$. Case a) corresponds to $x_o = 0.08$ and $\beta = 2.5$ for which there results $x_c \cong 0.098$, $c^{(b)}(x_o, \beta) \cong 0.211$, $f_c(x_o, \beta) \cong 0.246$ and $c^{(a)}(x_o, \beta) \cong 0.33$. Case b) corresponds to $x_o = 0.15$ and $\beta = 10$ for which there results $x_c \cong 0.171$, $c^{(b)}(x_o, \beta) \cong 0.106$, $f_c(x_o, \beta) \cong 0.119$ and $c^{(a)}(x_o, \beta) \cong 0.19$.




FIG. 2: Plot a) shows $c^{(a)}(x_o, \Gamma)$ as a function of β with $x_o = 0.1, 0.06, 0.024, 0.048$). Plot b) shows $c^{(a)}(x_o, \Gamma)$ as a function x_o for $\beta = 10, 20, 40, 60, 80, 100$.